

Methodological Module F-NW-def



**For the Design and Implementation
of Actions to Reduce GHG
Emissions from Deforestation
in Non-wetland Areas**

Version 1.1

Contents

Index of tables.....	4
Index of figures.....	4
1 Purpose and conditions of applicability	5
2 Segment delimitation	6
2.1 Activities considered in the segment.....	6
2.2 Historical, projection and results periods.....	6
2.3 Eligibility of areas and establishment of spatial limits.....	7
2.3.1 Collection of cartographic information.....	7
2.3.2 Map cross-referencing and coverage changes	9
2.3.3 Eligible areas	9
2.3.4 Compatibility with land use categories, land-use planning, and applicable environmental legislation.....	10
3 Identification and calculation of the segment's baseline scenario	11
3.1 Analysis of agents and causes of forest decline	11
3.1.1 Additional segment analysis factors	15
3.2 Considered carbon pools	17
3.3 GHG emission sources considered.....	18
3.4 GHG emission factors.....	19
3.5 Baseline scenario activity data.....	22
3.6 System and projection period.....	22
3.7 Estimated GHG emissions and removals from the baseline scenario	24
3.7.1 Sequence and segment calculations.....	24
4 Identification and calculation of the project scenario of the segment.....	26
4.1 Mitigation actions	26
4.2 Stratification of the project scenario	27
4.3 Carbon pools	27
4.4 GHG emission sources	27
4.4.1 Potential leakage area.....	27
4.5 GHG emission factors of the project scenario	28
4.6 Project scenario activity data.....	29
4.7 Estimated avoided emissions from the implementation of CCMP activities	29
4.8 Leakage estimation	30
5 Estimated net GHG emission reductions from the segment.....	31
6 Monitoring and quantifying results	32
6.1 Frequency of monitoring	32
6.2 Monitoring the implementation of the activities proposed for the segment.....	32
6.3 Emissions monitoring.....	32
6.4 Carbon stocks monitoring.....	33
6.5 Segment stratification.....	34
6.6 Monitoring of leakage management areas.....	34
6.7 Monitoring of increases in GHG emissions.....	35
6.8 Anthropogenic GHG emission reductions <i>ex-post</i> from the segment.....	35

6.9 Summary of results during the monitoring period	36
6.10 Monitoring and revalidation of the baseline scenario	37
6.11 Verifiable requirements in the implementation of the CCMP	37
6.12 Monitoring data and parameters	38
7 Document history.....	39

Index of tables

Table 1. Possible structure of the presentation of CCMP supporting information.	8
Table 2. Example of change or confusion matrix structure.	9
Table 3. Example of a structure for the presentation of eligibility information.	9
Table 4. Mapping indicators and data sources of the main activities associated with deforestation (drivers).	14
Table 5. Examples of actions to reduce deforestation and forest degradation in an indigenous reservation by improving local governance.	16
Table 6. List of cartographic inputs and sources used.	17
Table 7. Pools that can be included in the segment.	17
Table 8. GHG emission sources that can be included in the segment.	18
Table 9. Matrix of land use changes that may occur in the CCMP intervention area and in the F-NW-def segment.	20
Table 10. Leakage considered in a CCMP.	28
Table 11. Possible structure of the fire and burn occurrence reporting table.	33
Table 12. Process for calculating the segment's anthropogenic GHG <i>ex-post</i> emission reductions.	35
Table 13. Disaggregation of mitigation results obtained during the monitoring period.	37

Index of figures

Figure 1. Direct and underlying causes of deforestation.	13
Figure 2. Example of a projection period.	22

1 Purpose and conditions of applicability

This methodological module is designed for the implementation of climate change mitigation activities through actions to reduce deforestation in non-wetland forest areas and should be used in conjunction with Cercarbono modular methodologies that include the **F-NW-def** segment, which focuses on activities to **reduce GHG emissions from deforestation in non-wetland areas**.

The implemented mitigation reached through this type of activity and calculated with this module will be reported both independently for this module and in conjunction with the other segments implemented in a CCMP, which facilitates the correct assignment of the attributes of each credit type as well as the potential reporting of mitigation in the context of the NDCs in the corresponding land use category.

Scope of the methodological module

Land use category	Forest land.
Methodological category	Non-wetland forests.
Mitigation type	Reduction of GHG emissions.
Considered activities	Reduced deforestation.
Considered pools	Above-ground tree biomass, above-ground non-tree biomass, below-ground biomass, dead wood and coarse and fine litter, soil organic carbon.
Considered GHG emission sources	Biomass removal or burning by deforestation processes (including possible intermediate step of burning).

All areas of the CCMP belonging to this category form a separate segment. Recommendations are included for the delimitation of the segment, the identification and selection of baseline and project scenarios, relevant GHG emission sources and carbon pools, as well as for the quantification and monitoring of the results attained in the segment.

If a CCMP includes this **F-NW-def** segment, once validated, it cannot be excluded. The activity of this segment can be implemented in isolation or in conjunction with activities of other segments considered in the selected methodology in the same CCMP.

2 Segment delimitation

The delimitation of the segment requires the definition of the different elements that specify its scope:

- Time limits.
- Activities considered.
- Spatial limits.
- Carbon pools considered.
- Emission sources considered.

The **time limits** are defined at the CCMP level, for the total set of segments included in the CCMP. Other segment limits are defined in the following sections.

2.1 Activities considered in the segment

This module can be used for the implementation of activities focused on reducing GHG emissions through deforestation reduction processes in non-wetland forests.

2.2 Historical, projection and results periods

In addition to the time limits that are defined at the CCMP level, it is necessary to define for the segment the historical, projection and result periods, as detailed below. These periods should be the same for all deforestation reduction and degradation or land use change reduction segments included in the CCMP (segments F-NW-def, F-NW-deg, F-CW-def, F-CW-deg, F-CW-sfm, F-IW-Min-def, F-IW-Min-deg, F-IW-Orig-dra, NF-CW-mar, NF-CW-luc, NF-IW-Orig-dra and NF-IW-Min-luc, see **Annex 1 of the methodology** for identification and description).

- **Historical period (of historical emissions analysis)**¹: a period (in years) for which a trend in the agents and drivers of deforestation that is detectable in the activity data can be described and which is used to predict (estimate) the rate of deforestation that would occur during the projection period. This period should not be less than ten years.
- **Projection period**: is the time range (in years) for which projections are made in the baseline scenario based on the historical period. Emissions from deforestation are projected over this period. The starting year of this period should coincide with the project start date where the first CCMP interventions are carried out in the territory, covering the entire project duration or beyond.
- **Results period**: time range (in years) during which the activities of the segment and the results of these actions in terms of GHG emission reductions from deforestation of forest carbon content are monitored. The results period includes the verification periods during which the monitoring of GHG removals or GHG emission reductions is carried out. The duration of this period is equal to the duration of the CCMP.

¹ It should be aligned with the national or sub-national FREL/FRL.

2.3 Eligibility of areas and establishment of spatial limits

The areas that make up this segment:

- They must have been covered by forest for at least ten years prior to the start date of the activity. See the definition of forest in the most recent version of the ***Terms and Definitions of the Voluntary Certification Programme of Cercarbono***.
- They cannot be wetlands, as defined in the most recent version of the ***Terms and Definitions of the Voluntary Certification Programme of Cercarbono***.
- Areas with organic soils and areas with mineral soils are eligible.
- Implementation of mitigation activities in the segment must not cause disturbance to natural forests.
- They must be supported by ownership or administrative capacity by the CCMP.

This segment cannot overlap spatially with any segment established in another methodological module. The segment (portions of land where avoided deforestation actions will take place) must be explicitly identified, among others, to avoid double counting in the results. Once defined, the segment will be part of the CCMP area.

In grouping CCMPs, the spatial limits of the segment may change during implementation due to the inclusion of new instances. In this case, a validation of the new areas and an update of the mitigation potential calculations of the segment considering these areas will be necessary.

The segment should be explicitly delimited by considering its area and strata.

The strata are the potential forest **types**, grouped as having common characteristics for calculation purposes.

The eligibility analysis based on historical forest cover comprises the following steps:

2.3.1 Collection of cartographic information

Information sources from remote sensing, orthophotos, coverages or land-use planning tools developed by institutions in charge of official cartography in the country where the CCMP is implemented are allowed, which are classified in medium spatial resolution and high spatial resolution. Mapping with drones or Global Positioning System (GPS)² is also allowed, whose quality must be adequate and proportional to the scale of the project according to the guidelines defined by the ISO 19157:2013 Standard or by the institution in charge of the official cartography of the country where the CCMP is implemented.

² The use of free map viewers as a source of complementary information is allowed as long as the images or maps used are of the same date as the assessed period.

The mapping files must be compatible with a Geographic Information System in *shapefile* (.shp) format, in the coordinate system defined by the institution in charge of the official mapping of the country where the CCMP is implemented.

The minimum mappable area, understood as the minimum unit of interpretation of cartographic sources and corresponding to the working scale, must be equal to the minimum size established in the forest definition of the country where the CCMP is implemented.

The following spatial resolutions can be used:

- **Medium spatial resolution:** information with a spatial resolution of 30 m to 100 m, which allows defining working scales smaller than 1: 50,000, coming from spectral sensitivity systems or satellite images such as, for example, Landsat, SPOT, ALOS, AVNIR-2, ASTER and IRSS.
- **High spatial resolution:** information with a spatial resolution of less than 30 m, allowing working scales larger than 1: 50,000 to be defined, from spectrally sensitive systems, satellite, or aerial imagery, e.g., Sentinel, RapidEye, orthophotos and LiDar.

Consideration should be given to the final scale of the products and the relevance of the information sources to the size of the discrete areas included and the total CCMP area.

The use of cartographic information with date differences of no more than one year with respect to the dates of analysis is allowed.

The following table shows the structure of how the CCMP can present the information.

Table 1. Possible structure of the presentation of CCMP supporting information.

Department, state, or province	Geographical location or sub-scene detail	Type of data source*	Area of coverage (ha)	Date of data source

* Picture, orthophoto, official map or other applicable sources.

2.3.1.1 Information in raster format

The information in raster format should be used by specialised software for the interpretation of the images. Each process must be documented: pre-processing, corrections, enhancements, classification, assignment, and final interpretation of the images.

2.3.1.2 Information in vector format

The sources of information in vector format used must be identified, described, and supported. If vectorisation of images is required, the procedure used should be documented.

Whether raster or vector information is used, the procedure for arriving at the land cover maps at each of the dates of analysis must be supported.

2.3.1.3 Documentary information

The documentation used for the analysis must consider the entire area to be included in the segment and must support the existing coverage at the date for which the analysis is made. The cartographic interpretation must be supplemented to support the coverages at the date of commencement of the CCMP and those at the time of legal support.

2.3.2 Map cross-referencing and coverage changes

With the maps generated in each period, a change or confusion matrix must be created to show which coverages remain in forest and which become non-forest in the two periods of analysis.

A change or confusion matrix is a square matrix of $n \times n$, where n refers to the number of coverages. This matrix shows the relationship between two years of analysis corresponding to the area under study. The first row corresponds to the coverages determined in Year 1 (start date of the CCMP). The first column corresponds to the cover determined in Year 2 (ten years before the CCMP start date). The matrix includes the areas for each period of analysis to corroborate the accuracy of the classification and the changes in coverage to be considered within the eligibility analysis.

The following table shows how the CCMP **can** present the information for the change or confusion matrix.

Table 2. Example of change or confusion matrix structure.

Crossover Year1/Year2	Forest 1	Land cover 1-1	Land cover 1-2	Land cover 1-n	Total Year 1
Forest 2					
Land cover 2-1					
Land cover 2-2					
Land cover 2-n					
Total Year 2					

2.3.3 Eligible areas

Eligible areas must be determined according to cross-referenced information and presented in a traceable form, in **shapefile** (.shp) format where the information is evidenced and presented in a transparent manner.

The following table presents the structure of how the CCMP **can** present the information for eligible areas.

Table 3. Example of a structure for the presentation of eligibility information.

Item	Forest	Land cover LB1*	Land cover LB2*	Land cover LBn*	Total
Eligible area					

Item	Forest	Land cover LB1*	Land cover LB2*	Land cover LBn*	Total
Ineligible area					
Total (ha)					

* Land cover LB1: Land cover at the first time point; Land cover LB2: Land cover at the second time point; Land cover LBn: Land cover at the n -th time point.

2.3.4 Compatibility with land use categories, land-use planning, and applicable environmental legislation

The CCMP shall demonstrate the compatibility of the actions developed in the segment with the land use categories, if any, in the country where it is implemented.

If the initiative is intended to be implemented in environmentally protected areas, a permit or authorisation, as appropriate, must also be obtained from the environmental authority with jurisdiction in the area of intervention, which will verify the compatibility of the CCMP with the management instrument and the zoning established therein.

The CCMP must specify all applicable laws, statutes, and regulatory frameworks (local, regional, national, among others) in force in terms of categorisation or land management and must identify, implement, and periodically evaluate compliance with them.

Once the area eligibility analysis has been carried out, the spatial limits of the segment can be defined, selecting from the eligible areas those that will make up the segment.

3 Identification and calculation of the segment's baseline scenario

The baseline scenario in this module consists of estimating the amount of carbon in the pools and emissions by sources that would occur within the segment limits in the absence of the activities to be performed in the segment. The pools and GHG emission sources possible to consider in a CCMP are detailed in [Table 7](#) and [Table 8](#), respectively.

The calculation of GHG emissions of the baseline scenario is the result of the change in GHG emission sources and pools selected based on the change in the behaviour of the agents and drivers of deforestation, for which the following steps should be followed:

- Determine the location of the segment within the CCMP, adjusted to forest areas through the analysis of agents and drivers of deforestation ([Section 3.1](#)).
- Establish the historical period where activity data and GHG emission factors are measured for the calculation of historical emissions from deforestation ([Section 2.2](#)).
- Design and implement sampling for representative measurement of GHG emission factors ([Section 3.4](#)).
- Perform deforestation trend projection of the baseline scenario ([Section 3.6](#)).

3.1 Analysis of agents and causes of forest decline

The analysis of agents and drivers of deforestation starts from the preliminary analysis (see ***corresponding section in the methodology***) and is supported by secondary information collected on socio-economic variables of historical processes of deforestation and forest degradation. The agents and causes included are those that are associated with unsustainable uses of the forest, but also those that show the potential for sustainable management or leverage conservation processes including ethnic, cultural and livelihood conservation factors.

The analysis of the agents and causes of forest decline should be carried out in aggregate for all segments considered in [Section 2.1](#).

The analysis of agents and causes should be an iterative process as better and more up-to-date information becomes available to improve the effectiveness of CCMP actions. In its first iteration the main results should provide the territorial information inputs to generate:

- A first portfolio of REDD+ activities (a framework of possible activities is included in ***Annex 3 of the methodology***).
- The spatial delimitation of the segment areas.
- The temporal delimitation of the CCMP.
- The definition of the final location of the REDD+ activity segments.

It is recommended that the remaining iterations are carried out on an annual basis according to the circumstances of the CCMP. This means that the first diagnosis of causes and actors is done in the consolidation of the PDD. Once the first verification is done, one calendar year should be counted and the dialogues at the local level and the information on


socio-economic factors should be reprocessed to analyse the new behaviours of the agents and causes.

The CCMP should describe the agents and drivers of direct deforestation, as well as the associated underlying causes that will determine the dynamics of REDD+ activities (*Figure 1*). It is recommended to use a variety of information (e.g., expert consultation, participatory social assessments, literature review, etc.).

To this is added the knowledge of future conditions that directly or indirectly influence the decision of the different agents (e.g., new policies that encourage the production of a certain crop, policies around land use, among others).

Underlying causes are classified as those related to social, economic, demographic, technological, political, and institutional and cultural factors. The behaviour of the underlying and direct causes should be described at the project level.

Figure 1. Direct and underlying causes of deforestation.



Source: Adapted from Geist & Lambin (2002).

For the description of the agents and drivers of deforestation present in the segment, the combination of remotely sensed information with field-corroborated social dynamics data is recommended. For spatial analysis, mappable indicators associated with economic activities can be used. **Table 4** provides a framework of variables that can be considered and measured in an analysis of agents and drivers of deforestation.

The delimitation of the agents and drivers of deforestation analysis is based on the identification of micro-watersheds overlapping or adjacent to the project area or a smaller area, in

case a restricted distribution of such agents and drivers operating in the project area is demonstrated, which constitutes the reference area (see ***spatial limits section of the methodology***). The delimitation of micro-watersheds should follow the guidelines available at the national level (this methodology uses the micro-watershed as the unit of analysis, however, other similar elements that apply in each country can be integrated). In cases where micro-watersheds do not represent a logical unit of analysis of agents and drivers of deforestation (e.g., because there are external factors that influence these agents and drivers, such as administrative divisions or infrastructure elements that generate specific conditions in each sector), the CCMP may use, with due justification, a different spatial delimitation for the analysis of agents and drivers of deforestation.

Table 4. Mapping indicators and data sources of the main activities associated with deforestation (drivers).

Activity/driver of deforestation	Mapping indicator	Common data sources (national level)	Common data sources for GHG emissions estimation (national level)	Examples of other indirect data
Commercial agriculture	Large areas logged, post-harvest land use.	Historical satellite imagery (e.g., Landsat).	Traditional forest inventories / field measurements.	Commodity prices, agricultural censuses, share of gross domestic product, exports, among others.
Subsistence farming, smaller crops, and rotational crops	Small, logged areas, usually associated with rotation cycles.	Historical satellite images with high temporal density or high resolution to determine rotation pattern.	Traditional forest inventories / field measurements and targeted surveys.	Population growth in rural and urban areas, agricultural imports and exports, land use practices, among others.
Expansion of infrastructure	Road network, new mines, and built-up areas.	Historical satellite images.	Traditional forest inventories / field measurements.	Growth in urban and rural population, infrastructure development programmes, import and export prices of raw materials (mining).
Industrial or commercial harvesting of forest products	Small-scale canopy damage, logging roads and associated infrastructure.	Historical satellite imagery analysed in conjunction with concession areas. Direct analysis for recent years.	Traditional forest inventories / field measurements and harvest estimates from commercial forestry activities. GHG emission factors can be measured consistently over each historical period.	Rural and urban population growth, percentage of energy users and sources of energy, consumption patterns and their changes.

Activity/driver of deforestation	Mapping indicator	Common data sources (national level)	Common data sources for GHG emissions estimation (national level)	Examples of other indirect data
Extraction of forest products for subsistence, local and regional markets	Very small-scale canopy damage, understory impacts, footpaths.	<ul style="list-style-type: none"> - Limited historical data. - Information from local studies or national proxies. - Only long-term cumulative changes can be observed by satellite imagery. 	<ul style="list-style-type: none"> - Limited historical data. - Information from local scale studies. - Community-based monitoring has a key role. - Other indirect methods of measuring carbon stock changes can be employed. 	Land use practices (e.g., agricultural burning), links to other activity data attributable to burning, fire prevention and natural fires.
Other disturbances (e.g., uncontrolled fires)	Burn scars and associated impacts.	Historical fire-related satellite data, analysed in conjunction with Landsat-type data.	Regular estimation of emissions can be measured consistently for different periods depending on data availability.	

Source: Adapted from Kissinger *et al.*, 2012.

3.1.1 Additional segment analysis factors

In addition to the behaviour of the economic activities described above and summarised in **Table 4**, the following factors should be analysed in the segment:

Biophysical factors

Climate, soils, lithology, topography, relief, hydrology, and vegetation, which show spatial-temporal variation.

Economic and technological factors

Consider, for example, the commercialisation and growth of international timber markets or economic variables with low domestic costs (land, labour, fuel, etc.), increased product prices and the demands of remote urban and industrial centres.

Production factors

Analyse production systems and their influence on deforestation and forest degradation, whether they are in forest areas, legally or illegally established in the project reference area. For example: extractive industries, legal timber extraction, illegal timber extraction, cattle ranching, illicit crops, among others.

Demographic factors

The composition and distribution of the population, as well as the context in which the population interacts with other factors, are the most important demographic aspects for

understanding the pressure on land use and land cover changes, as well as the analysis of migration processes, which in turn are linked to other non-demographic factors, such as government policies, changes in consumption patterns and globalisation, which is clearly facilitated by the construction of infrastructure (e.g. access roads).

Institutional factors

Government policies play a major role in forest cover transformations, either directly or indirectly, mediating and interacting with demographic, economic, biophysical, and other factors. For example, access to land, capital, technology, and information are structured and often limited by national policies and institutions.

For the identification of the likely segments for the CSE, the available information on areas susceptible to restoration considered in national plans will be included in the analysis of agents and causes.

Territorial analysis

A product of the spatial information associated with the agents and causes is a map indicating how the different sources of pressure on the forest operate. This map should be easy to read and illustrative, as with this input it is recommended that participatory social mapping processes are carried out by means of a broad convocation of actors in the CCMP area. This process is achieved through the establishment of working groups in which it is confirmed whether what is detected in the mappable inputs is happening. This last step is what determines the diagnosis of the agents and causes of deforestation. It is also recommended to have as input the construction of timelines that include motivations, memories, histories, attitudes, values, perceptions, as well as personal and collective beliefs that affect decision-making.

With the socio-economic information compiled, a summary timeline of the factors that have generated the processes of deforestation and forest degradation must be constructed. In addition, correlations of events and trend analysis of these variables will be carried out for the most effective design of CCMP actions, a reference framework of actions is included in ***Annex 3 of the methodology***.

If, for example, the relationship between the analysis of agents and causes shows that the main agent of deforestation is the illegal occupants of extensions of land for the establishment of livestock in an indigenous reservation, and this is confirmed by the information on land use change, community testimonies and secondary information that describes historical processes of occupation of the reservation, after corroborating this process, actions should be generated from the CCMP such as those included in the table below.

Table 5. Examples of actions to reduce deforestation and forest degradation in an indigenous reservation by improving local governance.

Possible actions to reduce deforestation and forest degradation	
	Formulate and implement an ethnic-territorial planning instrument.

Administration measures	Strengthen the governance of indigenous people in their reservation through funding for their organisational structures and administrative capacity building for the design and implementation of projects.
Control measures	Implement a local early warning system for deforestation and forest degradation.
	Co-finance an agreement with the environmental authority to strengthen control processes in the reservation.
	Support the development of command-and-control measures, so that complaints about logging processes can be enforced without putting the community at risk.
Planning measures	Design and implement a roadmap for accessing financial mechanisms such as PES for forest cultural services.

One tool that can be included for the analysis of the current and future behaviour of the agents and drivers of deforestation is the construction of risk maps of forest loss, based on the variables analysed. If this alternative is implemented, the cartographic inputs and sources used must be traceable, for which it is recommended that **Table 6** be considered.

Table 6. List of cartographic inputs and sources used.

Mapping factor	Source	Variable that represents	Variable analysis	Data evaluation range	Criterium	Algorithm or equation used	Comments
ID	Filename	Units	Description				

In the framework of this methodology, risk maps are complementary tools for the analysis of agents and causes and, therefore, for the design of territorial actions to avoid deforestation or forest degradation; however, they do not replace projection systems and the inclusion of activities other than deforestation in the baseline scenario.

3.2 Considered carbon pools

The carbon pools included in the segment are those that can be measured to assess the carbon content in the baseline scenario and whose changes are assessed in the project scenario associated with the activities considered.

The pools that can be included in the segment are presented in **Table 7**.

Table 7. Pools that can be included in the segment.

Pool	Inclusion	Comments
Above-ground tree biomass	Yes	Pool subject to project activities. Includes woody tree biomass. Includes stems, stumps, branches, bark, seeds, and foliage. This pool is expected to be maintained due to forest management on forest land maintained as such.
Above-ground non-tree biomass	Opt.	Pool subject to project activities. Includes non-tree woody biomass and (shrubs and herbaceous). Includes stems, branches, bark, seeds, and foliage. This pool is expected to be maintained due to forest management on forest land maintained as such.

Pool	Inclusion	Comments
Below-ground biomass	Yes	Pool subject to project activities. Includes live root biomass greater than 2 mm in diameter. This pool is expected to be maintained due to forest management on forest land maintained as forest land.
Dead wood and coarse and fine litter	Opt.	Pool that may be subject to project activities in cases where it is identified as a key pool and monitoring is feasible or improved accuracy of measurement of its removals is considered. Includes non-living above ground wood, either standing or fallen such as dead roots and stumps greater than 10 cm in diameter.
Wood products	No	Can only be included in sustainable forest management activities (segments).
Soil organic carbon (SOC)	Opt.	Pool subject to project activities. Soil organic carbon content is expected to be maintained, due to avoided land cover changes (deforestation or forest degradation). Includes organic carbon from mineral and organic soils at a minimum depth of 30 cm and roots less than 2 mm in diameter.

3.3 GHG emission sources considered

The potential GHG emission sources included in the baseline scenario of this segment are due to deforestation and correspond accordingly to the GHG emissions avoided by these actions in the project scenario due to the implementation of the segment activities.

The GHG emission sources identified in the baseline scenario are to be monitored in the project scenario.

Table 8. GHG emission sources that can be included in the segment.

Process - source	GHG	Inclusion	Detail
Aquaculture (construction).	CO ₂	NA	Activity not considered in the segment.
Aquaculture (use) Emissions from aquaculture use.	N ₂ O	NA	Activity not considered in the segment.
Biomass removal or burning from deforestation processes (including possible intermediate step of burning).	CO ₂	Yes	Gas emitted by this source.
	CH ₄	Opt.	Gas emitted by this source.
	N ₂ O	Opt.	Gas emitted by this source.
Biomass removal or burning due to forest degradation processes (fragmentation).	CO ₂	NA	Gas emitted by this source.
	CH ₄	NA	Conservatively excluded.
	N ₂ O	NA	Conservatively excluded.
Biomass removal due to forest degradation processes (timber extraction).	CO ₂	NA	Gas emitted from this source.
	CH ₄	NA	Conservatively excluded.
	N ₂ O	NA	Conservatively excluded.
Conversion of forest land to other uses.	CO ₂	Yes	Gas emitted from this source.
Wood and fuelwood removal and charcoal production.	CO ₂	No	Activity not considered in the segment.
Conversion of other uses to forest land.	CO ₂	NA	Activity not considered in the segment.
Extraction - Excavation to allow construction of ports, harbours and marinas and filling or dredging to facilitate land elevation.	CO ₂	NA	Activity not considered in the segment.
Drainage - Agriculture, forestry, mosquito control.	CO ₂	NA	Activity not considered in the segment.

Process - source	GHG	Inclusion	Detail
Emissions and removals from drained inland organic soils on cropland, grassland, wetlands, settlements, and other land.	CO ₂	NA	Activity not considered in the segment.
Emissions from drainage ditches or channels.	CO ₂	NA	Activity not considered in the segment.
	CH ₄	NA	
Release of dissolved organic carbon (DOC) from organic soils to drainage water.	CO ₂	NA	Activity not considered in the segment.
Peat fires.	CO ₂	NA	Activity not considered in the segment.
	CH ₄	NA	
	CO	NA	
Emissions and removals from construction and operation of constructed wetlands for wastewater treatment (CW), including semi-natural treatment wetlands (SNTW).	CO ₂	NA	Activity not considered in the segment.
	CH ₄	NA	
	N ₂ O	NA	
Rewetting, revegetation, and creation - Conversion of drained to saturated soils through restoration of hydrology and re-establishment of vegetation. Re-vegetation on undrained soils.	CO ₂	NA	Activity not considered in the segment.
Rewetting, revegetation, and creation - Conversion of drained to saturated soils through restoration of hydrology and re-establishment of vegetation.	CO ₂	NA	Activity not considered in the segment.
Rewetted soils - CH ₄ emissions from change to natural vegetation following modifications to restore hydrology.	CH ₄	NA	Activity not considered in the segment.
Rewetting, revegetation, and creation - Re-vegetation of undrained soils.	CO ₂	NA	Activity not considered in the segment.
	CH ₄	NA	
Salt production (construction).	CO ₂	NA	Activity not considered in the segment.

In the case of burns and fires, following the IPCC (2006) guidelines, it is essential to understand the nature of these, to classify them as anthropogenic or different, and their calculation corresponds to the carbon fraction of the available fuel mass (biomass).

To make an estimate in a consistent manner, it is necessary:

- Obtain estimates of the area burnt.
- Estimate the mass of fuel available for combustion, which includes live biomass, litter, and dead wood.
- Select combustion factor.
- Select GHG emission factors.

3.4 GHG emission factors

GHG emission factors should be representative of the forest strata in the segment and should demonstrate internal consistency with the area where activity data is monitored and the project area.

Quantification should be performed on the pools affected by the significant sources (accumulating 90 % of carbon) and with measurement feasibility. For these pools, GHG emission factors are calculated based on forest inventories.

For field measurement it is recommended to follow the national forest inventory manuals, these inventories, and other processes of compiling information on GHG emission sources and carbon pools should have a representative number of samples to determine in each area, for each segment and for each stratum, the variables needed in the calculation of carbon content in all affected pools and for all selected sources.

To classify a source as significant, information on potentially significant GHG emission sources and their estimation is recorded, arranged in a table in ascending order of the total amount of carbon emitted in the historical period in the CCMP area, and all activities less than or equal to the 90th percentile are classified as significant. As definitive emission factors are often not available at the time of this calculation, they can be used from secondary information.

GHG emission factors are calculated on the pools that may be affected by the changes highlighted in green, as indicated in [Table 9](#) and net emissions are assumed (post deforestation land cover values are considered).

Above-ground and below-ground biomass pools should be included. In case any GHG emission source or carbon pool is not estimated, the reasons should be duly explained. Similarly, it is possible that information gaps may occur in the activity data. For these cases it is proposed to use the notation NA (not applicable) or NE (not estimated).

Field sampling (inventory) allows the compilation of data on forest structure and composition that feed allometric equations to estimate the carbon contained in the selected pools.

The rationale for the selection of allometric equations must be clearly documented. The selection of their parameters should be consistent with **Figure 1 of the methodology**. In case the CCMP does not have its own developments, it is recommended that this is done considering the process described in the reliability principle and a sub-criterion of taxonomic and ecological relevance, whereby equations are chosen according to their availability at the species, genus, family, or forest type scale, in that order of choice.

In [Table 9](#), the matrix includes possible changes in land use according to IPCC (2006); it is common for the CCMP to report changes in forest/non-forest categories. Both options are valid, if they are justified.

Table 9. Matrix of land use changes that may occur in the CCMP intervention area and in the F-NW-def segment.

Land use		x ₁ (ha)					Total (ha)
		Forest land	Agricultural land	Pasture	Settlements	Secondary vegetation	Other land
	Forest land	Degradation		Removals			

x ₂ (ha)	Agricultural land	Emissions						
	Pasture							
	Settlements							
	Secondary vegetation			Removals				
	Other land							
Total (ha)								

x: Represents the time variable.

x₁: Historical period.

x₂: Projection period.

 Transitions included in the methodology but not included in the segment.

 Transitions potentially included in the segment.

The selected allometric equations should be used in the data range in which they were constructed and follow the quantification recommendations of their authors (e.g., corrections for heteroscedasticity).

Individuals should be identified taxonomically, with herbarium support according to the allometric equations used and should preferably correspond to species scale. In case the allometric equations used are designed for ecosystems or forest types, identification of all species is not necessary.

Individuals not fully identified or without information at species level are recommended to be assigned the parameter values of the average by genus or family or the average for the species registered in each plot, in this order. In the absence of attributable data by taxonomic category, default data may be used as recommended in **Annex 4 of the methodology**.

The data in the field forms are evidence of monitoring and should be documented and available for verification and use in subsequent calculations.

For the calculation of net deforestation, the estimate of the statistically representative carbon content of the land covers that have replaced the forest will have to be included.

The **below-ground biomass** is assumed to degrade linearly, over a period of 20 years from the time of deforestation; therefore, the annual factor corresponds to 5 % of the total below-ground biomass of the forest concerned. These values are accounted for annually for 20 years, starting from the year after the deforestation/forest degradation. In the case of estimating emissions from deforestation in the **soil organic carbon** pool (optional to include), the carbon content is emitted in equal proportions over a period of oxidation (recommended twenty years) after the deforestation event occurs, so each annual estimate should include the expected portion of soil emissions for the year in which the estimate is made.

The emission factors calculated for this segment are the same for the baseline scenario and the project scenario.

3.5 Baseline scenario activity data

The selection of activities and the procedures for the calculation of activity data should be internally consistent with the baseline scenario. If new emission sources are identified, they should be included in the project scenario and the baseline scenario reassessed.

3.6 System and projection period


The choice of deforestation projection system should be a function of accuracy and relevance. To assess accuracy, the one that demonstrates the least error between model and actual data should be selected. Projections can be linear (trends or imputations of constant rates of deforestation), non-linear (e.g. logistic models) or models based on the probability of forest loss as a function of socio-economic or biophysical variables.

The appropriateness of the projection method is assessed in terms of choosing a reliable method (demonstrating its appropriateness through scientific references).

Figure 2 exemplifies the theoretical choice of method, where the total amount of emissions and potential GHG emission mitigation outcomes is the area under the curve (highlighted in blue).

The projection should include information for the historical period (annual emissions) that allows estimating the most realistic possible trend. In cases of linear trends, all annual data of the historical series should be included. For models that partially require information from the historical series (e.g., deforestation rate from a logistic model, calculated from two years), you should be conservative in the choice of method and years of the projection.

Figure 2. Example of a projection period.



In any scenario, the same FREL projection method and calculation steps and assumptions should be used in the overlap event. Using the same projection method includes:

- Using the same calculation equations, but verifying that each of the assumptions are met, e.g., if at the national level a deforestation rate is calculated using the two years with the lowest rates, at the project level the years with the lowest rates in the segment should be selected, not the same years as at the national level.
- Use the same period of historical data analysis.
- If there are assumptions that are not applicable to the segment, justify their non-inclusion.
- Verify that each of the applicable or non-applicable assumptions of the national level are met. It should be argued how they are excluded or adapted at the project level. For example, if the national baseline scenario excludes protected areas from the potential deforested forest, if there are no protected areas in the CCMP area, two options are available: either the exclusion criterion is justified as not applicable, or the areas that will not be deforested under land use, distribution or territorial planning figures not included in the national baseline scenario are identified.

A cut-off of the activity data monitored and during the historical period should be made. This will be the base information for the projection.

Possible equations for estimating the annual deforestation projection are averages, linear projections, or non-linear projections, such as logistic or models that correlate socio-economic and biophysical variables with the probability of deforestation. Two examples follow:

Logistic model:

$$AdefBL_x = \frac{Adef}{1 + e^{a+bx}} \quad (\text{Eq. 1})$$

Variable	Name	Units
$AdefBL_x$	Deforested area of the baseline scenario in year x (over the historical period).	ha
$Adef$	Area of forest susceptible to deforestation.	ha
e	Euler's constant.	
a	Model constant.	
bx	Annual deforestation rate for the last couple of years of the historical period.	

The imputation of a fixed annual deforestation rate, e.g., as proposed by Puyravaud (2003):

$$FDR = \left(\frac{1}{(x_2 - x_1)} \times \ln \frac{A_2}{A_1} \right) \cdot 100 \quad (\text{Eq. 2})$$

Variable	Name	Units
FDR	Fixed annual deforestation rate.	
x_1	Starting year of the period of analysis.	
x_2	Year-end of analysis period.	

Variable	Name	Units
A_1	Forest areas in the first year of the deforestation period analysed.	ha
A_2	Forest areas in the last year of the deforestation period analysed.	ha

In this case, the annual deforested area of the baseline scenario of the deforestation segment (A_{defBL_t}) would be calculated as:

$$A_{defBL_t} = FDR \times A_{def} \quad (\text{Eq. 3})$$

Variable	Name	Unit
A_{defBL_t}	Annual deforested area of the baseline scenario of the deforestation segment.	ha
FDR	Fixed annual deforestation rate.	
A_{def}	Area of forest susceptible to deforestation.	ha

3.7 Estimated GHG emissions and removals from the baseline scenario

The total GHG emissions and removals of the baseline scenario is the sum of the annual emissions of the projection period from deforestation in the segment.

3.7.1 Sequence and segment calculations

Process	Variable and calculation	Data source
Baseline scenario (estimated future GHG emissions in the absence of the project)		
Analysis of agents and drivers of deforestation.		Done by the developer.
Temporal delimitation.	t = CCMP year index. T = Total CCMP duration, in years.	Defined by developer.
Area delimitation.		
Reference area.		GIS layers defined by the developer based on the possibilities and analysis of actors and causes.
Potential leakage area.		
Leakage management area.		
Define the potential deforestation forest segment.		
The stratum index of the segment's baseline scenario is defined.	f	Defined in the methodology.
Define the total number of strata of the segment baseline scenario.	TSSBL	Defined by the developer according to the characteristics of the forest.
Define the area of each stratum f of the segment baseline scenario.	$ASBL_f$	
Determine the above-ground biomass per unit area of each stratum f of the segment's baseline scenario.	AbS_f	Inventories or acceptable references.

Process	Variable and calculation	Data source
Determine the below-ground biomass per unit area of each stratum f of the segment baseline scenario.	BbS_f	Field measurement or allometric model supported.
Determine the dead wood and litter per unit area of each stratum f of the segment baseline scenario.	DwS_f	Acceptable inventories or references.
Determine soil organic carbon per unit area of each stratum f of the segment's baseline scenario.	$SocS_f$	Field measurement or acceptable references.
Define emission sources.		CO ₂ , CH ₄ and N ₂ O, as selected.
Calculate emission factors for above-ground biomass and dead wood and litter (if included) for each stratum f of the segment baseline scenario.	$EF S_f = AbS_f + DwS_f$ (Eq. 4)	Calculation.
Calculate annual below-ground biomass emission factors for each stratum f of the segment baseline scenario.	$EF BbS_f = \frac{BbS_f}{20}$ until $t = 20$. (Eq. 5)	Calculation.
Calculate annual soil organic carbon emission factors for each stratum f of the segment's baseline scenario.	$EF SocS_f = \frac{SocS_f}{20}$ until $t = 20$. (Eq. 6)	Calculation.
Estimate annual deforestation activity data for each stratum f of the segment's baseline scenario.	$ADSBL_{t,f}$	Projection based on analysis of agents and drivers of deforestation.
Estimate deforestation emissions in each year t and each stratum f of the segment baseline scenario.	$CO2ESBL_{t,f} = ADSBL_{t,f} * (EF S_f + EF BbS_f + EF SocS_f)$ (Eq. 7)	Calculation.
Estimate cumulative emissions from deforestation in all strata of the segment baseline scenario.	$CO2ESBL = \sum_{t=1}^T \sum_{f=1}^{TSSBL} CO2ESBL_{t,f}$ (Eq. 8)	Calculation.

4 Identification and calculation of the project scenario of the segment

The project scenario depends mainly on the activities carried out on the territory and their effect on deforestation. The calculation of GHG emissions should correspond to the result of the change in GHG emission sources and the selected carbon pools for which the following general steps should be followed:

- Calculation of total and annual emissions of the baseline scenario expected for the projection period.
- Implementation of project actions in the territory.
- Quantification of results.

In addition, leakage and compliance with safeguards and Sustainable Development Goals (SDGs) should be analysed and described jointly for all segments considered in the CCMP.

GHG emission reductions directly resulting from non-REDD+ activities implemented as part of the CCMP (e.g., from the implementation of efficient cookstoves) shall be excluded from the calculations and only their indirect effects on the considered segments shall be considered.

4.1 Mitigation actions

Once the segment activity data has been estimated and the agents and drivers of deforestation in the segment have been identified, the actions to be implemented in the segment are determined, initiated, and documented.

In the implementation of the actions considered in this methodology, actions should be territorial, i.e., not exclusively focused on policy changes, although if they exist, they should be reported.

Territorial actions must be supported by the commitment of landowners and landholders to contribute to the project's actions.

Actions can be developed exclusively with the communities, in alliances with government institutions or with private actors. In each case, the starting date of the activity and the period in which deforestation reduction is generated due to its implementation should be specified.

Note 1: The longest range of time that any project activity is generating change in any project activity (in any segment) determines the duration of the CCMP.

Note 2: The time at which the generation of changes in REDD+ activities by CCMP activities begins determines the end point of the historical period and the beginning of the projection period (without and with project scenarios).

Actions to reduce deforestation should be aligned with the current forest policy in the country implementing the CCMP. Possible actions to reduce deforestation at project level are included in **Annex 3 of the methodology**.

Actions to reduce deforestation can be synergistic, complementary, or identical to those undertaken to achieve avoided forest degradation.

The avoided deforestation of the project scenario arises from the comparison of the expected deforestation of the corresponding segment and the annual data during the result period.

4.2 Stratification of the project scenario

When the segment has different stratification criteria or classes in the stratification criteria compared to the baseline scenario, it will be necessary to stratify differently from the baseline scenario.

As in the baseline scenario, in any of the cases where stratification is necessary, it will be necessary to define the land cover of each stratum in each segment. If, in any of the segments, scenarios or in the implementation of activities, no subdivision of areas is required, a single stratum will be considered to exist (and therefore the corresponding sub-index will have a single value equal to one).

4.3 Carbon pools

All pools included in the baseline scenario of the segment must be considered in the project scenario (following the principle of internal consistency). Furthermore, no pools may be added or removed during the duration of the CCMP.

In the framework of this methodology, the inclusion of at least the above-ground and below-ground biomass pools in the segment is mandatory.

All selected pools are considered to remain constant in areas that remain as forest and therefore the values defined for each pool remain static for the duration of the CCMP.

4.4 GHG emission sources

The GHG emission sources that can be included or excluded from the project activity are shown in **Table 8**. Their selection should demonstrate internal consistency with the emission sources included in the baseline scenario.

4.4.1 Potential leakage area

Based on the characterisation of agents and drivers of deforestation and forest degradation, a potential leakage area, outside the CCMP monitoring area, is defined based on four criteria:

1. Areas where the same productive activities associated with the agents and drivers of deforestation or forest degradation are present.
2. Ecosystem equivalence with the project area.
3. Micro-watersheds adjacent to the project monitoring area (if in the reference area).
4. Areas of stable forest.

Where demonstrable evidence can be collected that deforestation in the potential leakage area is attributable to deforestation agents that are not linked to the CCMP area, the detected deforestation is not attributed to the project activity and will not be considered as leakage.

Over this potential leakage area, a leakage management area (defined in the ***spatial limits section of the methodology***) is delimited.

The possible sources of GHG emissions due to leakage that can be included or excluded from the project activity are shown in ***Table 10***.

Table 10. Leakage considered in a CCMP.

Source	GHG	Inclusion	Explanation
Displacement by grazing and live-stock production	CO ₂	Optional	To be considered if significant in the potential leakage area.
	CH ₄	Optional	To be considered if significant in the potential leakage area.
	N ₂ O	No	Excluded (manure management is not included in the scope of this methodology).
Displacement by agricultural activities	CO ₂	Optional	To be considered if significant in the potential leakage area.
	CH ₄	Optional	To be considered if significant in the potential leakage area.
	N ₂ O	No	Excluded.
Increased fertiliser use	CO ₂	No	Excluded.
	CH ₄	No	Excluded.
	N ₂ O	Optional	To be considered if significant in the potential leakage area.
Timber harvesting	CO ₂	Optional	To be considered if significant in the potential leakage area.
	CH ₄	Optional	To be considered if significant in the potential leakage area.
	N ₂ O	No	Excluded.
Deforestation	CO ₂	Yes	To be considered if significant in the potential leakage area.
	CH ₄	Optional	To be considered if significant in the potential leakage area.
	N ₂ O	No	Excluded.

The relevance of the inclusion of leakage in the CCMP is defined by entering the monitoring area in the CCMP. If the project is in overlap with a reference level, leakage is not accounted for.

4.5 GHG emission factors of the project scenario

All GHG emission factors included by the CCMP in the baseline scenario must be considered in the project scenario.

4.6 Project scenario activity data

The selection of activities and the procedures for the calculation of activity data should be internally consistent with the baseline scenario. If new emission sources are identified, they should be included in the project scenario and the baseline scenario reassessed.

Monitoring of the segment activity data must be performed annually and over the years of the results period, both in the segment area, the monitoring area, and the leakage area.

4.7 Estimated avoided emissions from the implementation of CCMP activities

The sequence and calculations for segment **Reducing GHG emissions from deforestation in non-wetland areas (F-NW-def)**, are summarised below.

Annex 4 of the methodology lists different sources of complementary information useful for the estimation and calculation of some variables mentioned below.

Process	Variable and calculation	Data source
Project scenario (estimated future GHG emissions if the project were to be implemented)		
Define actions to reduce deforestation in the segment.		Defined by the developer.
Define the stratum index of the segment's project scenario deforestation.	g	Defined by the methodology.
Define the total number of strata of the segment project scenario.	TSSP	Defined by the developer, depending on the characteristics of the forest.
Define the area of each stratum g of the segment project scenario.	ASP_g	Determined by the developer.
Estimate the annual deforestation activity data for each stratum g of the segment project scenario.	$ADSP_{t,g}$	Projection based on overlap condition with a FREL or agent and cause analysis and effectiveness of planned activities.
Determine the above-ground biomass per unit area of each stratum g of the segment project scenario.	AbS_g	Acceptable inventories or references.
Determine the below-ground biomass per unit area of each stratum g of the segment project scenario.	BbS_g	Field measurement or allometric model supported.

Process	Variable and calculation	Data source
Determine the dead wood and litter per unit area of each stratum g of the segment project scenario.	DwS_g	Acceptable inventories or references.
Determine soil organic carbon per unit area of each stratum g of the segment project scenario.	$SocS_g$	Field measurement or acceptable references.
Calculate the above-ground biomass and dead wood and debris (if included) emission factors for deforestation for each stratum g of the segment project scenario.	$EFS_g = AbS_g + DwS_g$ (Eq. 9)	Calculation.
Calculate annual below-ground biomass emission factors for each stratum g of the segment project scenario.	$EFBbS_g = \frac{BbS_g}{20}$ until $t = 20$. (Eq. 10)	Calculation.
Calculate annual soil organic carbon emission factors for each stratum g of the segment project scenario.	$EFSocS_g = \frac{SocS_g}{20}$ until $t = 20$. (Eq. 11)	Calculation.
Calculate the deforestation emissions in each year t and each stratum g of the segment project scenario.	$CO2ESP_{t,g} = ADSP_{t,g} * (EFS_g + EFBbS_g + EFSocS_g)$ (Eq. 12)	Calculation.
Calculate deforestation emissions for all strata of the segment project scenario.	$CO2ESP = \sum_{t=1}^T \sum_{g=1}^{TSSP} CO2ESP_{t,g}$ (Eq. 13)	Calculation.

4.8 Leakage estimation

For the potential leakage area, estimates of potential leakage emissions resulting from the implementation of the activities of all segments considered in the CCMP should be made as indicated in the methodology.

5 Estimated net GHG emission reductions from the segment

Using the same method as for estimating activity data and emission factors, mitigation results are calculated annually, comparing the expected baseline scenario data with that obtained because of the implementation of the segment's actions.

Based on **Table 9** and **Figure 2**, the point at which the historical period ends, and the results period (projection period) begins is identified, which corresponds to the point at which the segment's actions influence deforestation.

Using the same methods and procedures used to estimate activity data and emission factors in the historical period, during the accreditation period, the volume of results is calculated for each year by comparing the expected (projected) data with that obtained because of the implementation of the segment's actions.

The results are expressed annually in tonnes of carbon dioxide (tCO₂) over the entire accreditation period.

If a CCMP overlaps with a FREL that includes national circumstances, these will be subject to methodological reconstruction and therefore quantifiable at the project level, provided that the project area meets the assumptions underpinning the assignment of such circumstances.

Process	Variable and calculation	Data source
Estimation of the total mitigation potential of the deforestation segment		
Calculate the total mitigation in each year t of the segment.	$TMS_t = \sum_{f=1}^{TSSBL} CO2ESBL_{t,f} - \sum_{g=1}^{TSSP} CO2ESP_{t,g}$ <p>(Eq. 14)</p>	Calculation.
Calculate total segment mitigation.	$TMS = \sum_{t=1}^T TMS_t$ <p>(Eq. 15)</p>	Calculation.

6 Monitoring and quantifying results

6.1 Frequency of monitoring

In this segment, monitoring of GHG emission reductions should be annual, while monitoring of GHG emissions should be done on a continuous basis, as outlined in [Section 6.3](#).

Activity data should be monitored during the years of the results period, in the CCMP area and in the potential leakage and leakage management areas.

6.2 Monitoring the implementation of the activities proposed for the segment

The CCMP activities implemented within the segment area must be consistent with the project area management plans and the PDD. The CCMP shall include, in the monitoring report, a summary of the activities carried out in the segment during each verification period and their effectiveness in terms of climate change mitigation, in the context of the activities proposed in the PDD, comparing what was planned with what was implemented.

In the segment of reducing GHG emissions from deforestation in non-wetland areas, activity data should be monitored annually over the years of the results period, in the segment area and in the potential leakage and leakage management areas (which is defined at the project level and not at the segment level).

If the data used for the emission factors applicable to the segment come from the host country's FREL, the CCMP shall use these factors. If a more recent version of the FREL defines updated values, the CCMP shall update the calculations in the verifications using these updated values.

If the data used for the emission factors applicable to the segment come from forest inventories and it is evident that there are no significant changes in emission factors in the monitored categories from the baseline scenario, the emission factors used in the project scenario should be the same and a repeat forest inventory is not necessary.

6.3 Emissions monitoring

This module only covers non-CO₂ gas emissions from forest fires. Increases in these emissions should only be estimated and accounted for if they were included in the baseline scenario.

To estimate the increase in GHG emissions due to forest fires in the potential leakage area, it should be assumed that forest clearing is done by burning the forest. The parameter values used to estimate emissions shall be the same as those used to estimate forest fires in the baseline scenario, except for the initial carbon buffers, which shall be those of the initial forest classes burned in the potential leakage area.

The CCMP shall keep a logbook of burn and fire occurrence, where the information shown in [Table 11](#) shall be reported. Based on this table, and according to the procedures

established in the CDM tool³, GHG emissions will be estimated for each occurrence and then the annual sum and for the corresponding verification periods.

Table 11. Possible structure of the fire and burn occurrence reporting table.

Date	Stratum	Affected area (ha)	Burnt biomass (%)	Comments

The result of the estimates should be reported using the same table formats used in the *ex-ante* assessment of baseline GHG emissions from forest fires in the CCMP area.

6.4 Carbon stocks monitoring

To quantify the actual GHG removals and GHG emission reductions achieved in the segment, it is necessary to monitor the changes in carbon stocks and GHG emissions within the segment by monitoring the following components:

- **Land use and land cover change within the segment.** It is necessary to monitor all forest areas that are converted to non-forest areas. Monitoring results must be presented in *ex-post* tables of activity data by stratum. It is mandatory that this monitoring is carried out throughout the duration of the CCMP.
- **Changes in carbon stocks.** In most cases, emission factors per land use or land cover category will not change during a fixed reference period and will not need to be monitored. However, monitoring of carbon buffers is mandatory within the segment for areas subject to a significant decrease in project scenario carbon buffers according to the initial assessment. These will be areas subject to controlled deforestation and planned harvesting activities, such as logging, fuelwood collection and charcoal production. In these areas, changes in carbon buffers should be estimated at least once after each harvesting event.
- **Impacts of natural disturbances and other catastrophic events.** Decreases in carbon stocks and increases in GHG emissions are subject to monitoring and must be accounted for under the project scenario, where significant, even if such decreases are due to natural causes, e.g. in case of forest fires or natural disturbances such as hurricanes, earthquakes, volcanic eruptions, tsunamis, floods, droughts and the like, or human-induced events, including those over which the project proponent has no control (such as fires, acts of terrorism and war). In the case of forest fires, non-CO₂ emissions from forest fires should also be accounted for.
- **Estimated total changes in actual net carbon buffers and emissions (including leakage) of GHGs in the segment area.** Considering the above elements, the estimated total changes of actual net carbon buffers and GHG emissions in the segment should be calculated and summarised in a table.

³ [A/R Methodological Tool](#): Estimation of non-CO₂ GHG emissions resulting from burning of biomass attributable to an A/R CDM project activity (Version 04.0.0).

6.5 Segment stratification

As in the case of the project scenario, when different stratification criteria or classes in the stratification criteria are present during the implementation of the activities compared to the baseline scenario, a stratification different from the baseline scenario (and possibly also different from the project scenario) will be necessary.

As in the baseline and project scenarios, in any case where stratification is necessary, the land cover of each stratum in each segment must be defined. In case no subdivision of areas is required in any of the segments, scenarios or in the implementation of activities, a single stratum will be considered to exist (and therefore the corresponding sub-index will have a single value equal to one).

6.6 Monitoring of leakage management areas

The potential leakage area is an area covered by forest at the start of the CCMP, where the same agents and drivers that generate emissions from REDD+ activities in the project area may operate, while the leakage management area is a precise limit where leakage action has been identified and must be controlled. Evidence for identifying leakage includes:

- Deforestation processes outside the avoided deforestation segment.
- Forest degradation processes outside the avoided forest degradation segment.
- Displacement of livestock or land grabbing activities.
- Displacement of other productive activities associated with deforestation or forest degradation.

This area is monitored for activity data to be determined using the same methods applied to monitor deforestation activity data in the segment.

In the operation of the project and as a product of the monitoring and management of the information, a process of constant control of leakage must be established, which includes:

- A geographical delimitation of the areas where the control is carried out.
- Changes in carbon buffers and GHG emissions associated with leakage prevention activities.
- The decrease in carbon buffers and increase in GHG emissions due to leakage displacement activity.

Based on the above elements, it is necessary to calculate the total estimated actual leakage associated with the CCMP, in aggregate for all segments considered.

In the case of CCMPs not fully overlapping with a FREL, increases in deforestation in the leakage management area, after a control process, will be deducted from the project accounting at the next verification. In the case of overlap with a FREL, no discounts are made in the accounting, but actions for leakage reduction are formulated from the project.

The result of the *ex-post* estimates of carbon stock changes should be reported using the same table formats used in the *ex-ante* assessment of the baseline carbon stock changes in the potential leakage area.

The verifying VVB shall determine whether the documentation provided by the CCMP proponent represents sufficient evidence to consider that the detected deforestation is not attributable to the project activity and therefore not leakage.

6.7 Monitoring of increases in GHG emissions

Increases should only be estimated and accounted for if GHG emissions from forest fires (burning) are included in the baseline scenario.

To estimate the increase in GHG emissions due to forest fires in the potential leakage area, it should be assumed that forest clearing is done by burning the forest. The parameter values used to estimate emissions will be the same as those used to estimate forest fires in the baseline scenario, except for the initial carbon buffers, which will be those of the initial forest classes burned in the potential leakage area.

The result of the estimates shall be reported using the same table formats used in the *ex-ante* assessment of baseline GHG emissions from forest fires in the segment.

6.8 Anthropogenic GHG emission reductions *ex-post* from the segment

Ex-post estimated net anthropogenic GHG emissions should be reported using the same table format used for the *ex-ante* assessment. **Annex 4 of the methodology** lists different sources of supplementary information useful for the estimation and calculation of some variables which are mentioned below, together with relevant considerations, in **Table 12**.

Table 12. Process for calculating the segment's anthropogenic GHG *ex-post* emission reductions.

Process	Variable and calculation	Data source
Monitoring and quantification of results (calculation of the reductions reached in the segment)		
Index of the effectively implemented stratum of the deforestation segment.	h	Defined by the methodology.
Define the total number of strata effectively existing in the segment.	TSSE	Defined by the developer based on the characteristics of the forest.
Define the area of each stratum h effectively existing in the segment.	ASE_h	Determined by the developer.

Process	Variable and calculation	Data source
Monitor and determine the areas effectively deforested in each year t and each stratum h in the segment.	$ADSE_{t,h}$	Monitoring.
Monitor and determine the emissions from deforestation leakage that effectively occurred in each year t in the potential leakage area.	$EDLE_t$	Monitoring and calculation.
Define the number of years from the start of the project until the time of monitoring corresponding to reporting period x (where x is the ordinal of the reporting period).	Tx	Defined by developer.
Estimate emissions from deforestation in all strata during monitoring period x of the segment's baseline scenario.	$CO2ESBL_x = \sum_{t=1}^{Tx} \sum_{f=1}^{TSSBL} CO2ESBL_{t,f}$ (Eq. 16)	Calculation.
Quantify emissions occurring in all strata of the segment during monitoring period x .	$CO2ESE_x = \sum_{t=1}^{Tx} \sum_{h=1}^{TSSE} \sum_{f=1}^{TSSBL} \left(ADSE_{t,h} \right. \\ \left. * (EFS_f + EFBbS_f + EFSocS_f) \right)$ (Eq. 17)	Calculation.
Calculation of mitigation reached during the reporting period		
Quantify segment mitigation during reporting period x .	$SM_x = CO2ESBL_x - CO2ESE_x - SM_{(x-1)} *$ (Eq. 18)	Calculation.

* Where $SM_{(x-1)}$ is the segment mitigation during the previous reporting period.

Variable	Name	Units
SM_x	Mitigation reached in the segment during reporting period x .	tCO ₂

6.9 Summary of results during the monitoring period

The annual mitigation achieved in the segment during reporting period x (SM_x) should be disaggregated together with the carbon buffer and presented according to [Table 13](#). The results for each monitoring year should be presented in separate rows.

Table 13. Disaggregation of mitigation results obtained during the monitoring period.

Segment:													
Year	Baseline scenario (t-CO ₂ e)						Monitoring period results (t-CO ₂ e)						Total gross reductions (t-CO ₂ e)
	Carbon stocks				Emissions		Carbon stocks				Emissions		
	Above ground biomass	Below ground biomass	Dead wood and litter	Soil organic carbon	N ₂ O	CH ₄	Above ground biomass	Below ground biomass	Dead wood and litter	Soil organic carbon	N ₂ O	CH ₄	
Total													

6.10 Monitoring and revalidation of the baseline scenario

Baseline scenarios, regardless of the approach chosen to establish them, need to be revised over time because the agents, drivers and underlying causes of deforestation change dynamically. They also need to be updated when new instances are added to grouping CCMPs.

Frequent and unplanned updating of the baseline scenario can create serious market uncertainties. Therefore, the baseline scenario should be reviewed every five years, choosing historical and projection periods that do not create inconsistencies and inconsistencies with the periods already verified and thus with the results obtained and accredited. Where an applicable jurisdictional, subnational, or national baseline scenario is available, baseline scenarios may be reassessed earlier, according to the elements below. The tasks involved in revising the baseline scenario are:

- Update information on agents, drivers, and underlying causes of deforestation.
- Periodically collect information on the agents, drivers, and underlying causes of deforestation in the reference area as these are essential to improve future deforestation projections and project activity design. Information should be collected that is relevant to understanding the agents of deforestation, drivers, and underlying causes. When a spatial model is used to locate future deforestation, new data on the spatial driving variables used to model deforestation risk should be collected as they become available. They should be used to create updated spatial datasets and new "Driver Maps" for the subsequent fixed reference period.
- Adjust the land use and land cover change component of the baseline scenario.
- Adjust the annual reference deforestation areas.
- Adjust the location of projected reference deforestation.
- Adjust the carbon component of the baseline scenario.

6.11 Verifiable requirements in the implementation of the CCMP

The calculations of emission factors, activity data, historical period and projection method are performed by means of a verifiable methodological reconstruction, based on the execution of the baseline and project scenario building steps of the methodology in the segment, and consistent with the official procedures reported in the FRELS. In case this reference is not available in the country, other academically based procedures can be followed.

In line with the principle of transparency, all information necessary for the reconstruction of the segment results should be documented.

6.12 Monitoring data and parameters

The data and parameters to be monitored are presented in *Table 11* and *Table 12*.

7 Document history

Version	Date	Comments or changes
1.0	03.11.2022	Initial version of the document is open for public consultation from 03.11.2022 to 02.12.2022.
1.1	23.02.2023	Final version with integrated comments from the public consultation.